Topic #9

Miscellaneous

BC only

AP® CALCULUS BC 2002 SCORING GUIDELINES

Question 6

The Maclaurin series for the function f is given by

$$f(x) = \sum_{n=0}^{\infty} \frac{(2x)^{n+1}}{n+1} = 2x + \frac{4x^2}{2} + \frac{8x^3}{3} + \frac{16x^4}{4} + \cdots + \frac{(2x)^{n+1}}{n+1} + \cdots$$

on its interval of convergence.

(a) Find the interval of convergence of the Maclaurin series for f. Justify your answer.

(b) Find the first four terms and the general term for the Maclaurin series for $f'(x)$.

(c) Use the Maclaurin series you found in part (b) to find the value of $f'\left(-\frac{1}{3}\right)$.



3. The figure above shows the graphs of the line $x = \frac{5}{3}y$ and the curve C given by $x = \sqrt{1 + y^2}$. Let S be the shaded region bounded by the two graphs and the x-axis. The line and the curve intersect at point P.

(a) Find the coordinates of point P and the value of $\frac{dx}{dy}$ for curve C at point P.

(b) Set up and evaluate an integral expression with respect to y that gives the area of S.

(c) Curve C is a part of the curve $x^2 - y^2 = 1$. Show that $x^2 - y^2 = 1$ can be written as the polar equation $r^2 = \frac{1}{\cos^2\theta - \sin^2\theta}$.

(d) Use the polar equation given in part (c) to set up an integral expression with respect to the polar angle θ that represents the area of S.

$\frac{dx}{dy} = \frac{1}{2}(1 + y^2)^{-1/2}(2y)$

$\frac{dx}{dy} = \frac{y}{\sqrt{1+y^2}} = \frac{y}{x}$

END OF PART A OF SECTION II

$\int r$

$x^2 - y^2 = 1$

$r^2\cos^2\theta - r^2\sin^2\theta = 1$

$r^2(\cos^2\theta - \sin^2\theta = 1$

$r^2 =$



2. The curve above is drawn in the xy-plane and is described by the equation in polar coordinates $r = \theta + \sin(2\theta)$ for $0 \le \theta \le \pi$, where r is measured in meters and θ is measured in radians. The derivative of r with respect to θ is given by $\frac{dr}{d\theta} = 1 + 2\cos(2\theta)$.

   (a) Find the area bounded by the curve and the x-axis.

   (b) Find the angle θ that corresponds to the point on the curve with x-coordinate -2.

   (c) For $\frac{\pi}{3} < \theta < \frac{2\pi}{3}$, $\frac{dr}{d\theta}$ is negative. What does this fact say about r? What does this fact say about the curve?

   (d) Find the value of θ in the interval $0 \le \theta \le \frac{\pi}{2}$ that corresponds to the point on the curve in the first quadrant with greatest distance from the origin. Justify your answer.

WRITE ALL WORK IN THE TEST BOOKLET.

3. An object moving along a curve in the xy-plane is at position $(x(t), y(t))$ at time t, where

$$\frac{dx}{dt} = \sin^{-1}\left(1 - 2e^{-t}\right) \text{ and } \frac{dy}{dt} = \frac{4t}{1 + t^3}$$

for $t \geq 0$. At time $t = 2$, the object is at the point $(6, -3)$. (Note: $\sin^{-1} x = \arcsin x$)

(a) Find the acceleration vector and the speed of the object at time $t = 2$.

(b) The curve has a vertical tangent line at one point. At what time t is the object at this point?

(c) Let $m(t)$ denote the slope of the line tangent to the curve at the point $(x(t), y(t))$. Write an expression for $m(t)$ in terms of t and use it to evaluate $\lim_{t \to \infty} m(t)$.

(d) The graph of the curve has a horizontal asymptote $y = c$. Write, but do not evaluate, an expression involving an improper integral that represents this value c.

WRITE ALL WORK IN THE PINK EXAM BOOKLET.

END OF PART A OF SECTION II

5. Consider the differential equation $\frac{dy}{dx} = 5x^2 - \frac{6}{y-2}$ for $y \neq 2$. Let $y = f(x)$ be the particular solution to this differential equation with the initial condition $f(-1) = -4$.

   (a) Evaluate $\frac{dy}{dx}$ and $\frac{d^2y}{dx^2}$ at $(-1, -4)$.

   (b) Is it possible for the x-axis to be tangent to the graph of f at some point? Explain why or why not.

   (c) Find the second-degree Taylor polynomial for f about $x = -1$.

   (d) Use Euler's method, starting at $x = -1$ with two steps of equal size, to approximate $f(0)$. Show the work that leads to your answer.



3. The graphs of the polar curves $r = 2$ and $r = 3 + 2\cos\theta$ are shown in the figure above. The curves intersect when $\theta = \frac{2\pi}{3}$ and $\theta = \frac{4\pi}{3}$.

   (a) Let R be the region that is inside the graph of $r = 2$ and also inside the graph of $r = 3 + 2\cos\theta$, as shaded in the figure above. Find the area of R.

   (b) A particle moving with nonzero velocity along the polar curve given by $r = 3 + 2\cos\theta$ has position $(x(t), y(t))$ at time t, with $\theta = 0$ when $t = 0$. This particle moves along the curve so that $\frac{dr}{dt} = \frac{dr}{d\theta}$. Find the value of $\frac{dr}{dt}$ at $\theta = \frac{\pi}{3}$ and interpret your answer in terms of the motion of the particle.

   (c) For the particle described in part (b), $\frac{dy}{dt} = \frac{dy}{d\theta}$. Find the value of $\frac{dy}{dt}$ at $\theta = \frac{\pi}{3}$ and interpret your answer in terms of the motion of the particle.

WRITE ALL WORK IN THE PINK EXAM BOOKLET.

END OF PART A OF SECTION II

CALCULUS BC
SECTION II, Part B
Time—45 minutes
Number of problems—3

No calculator is allowed for these problems.

4. Consider the differential equation $\frac{dy}{dx} = 6x^2 - x^2y$. Let $y = f(x)$ be a particular solution to this differential equation with the initial condition $f(-1) = 2$.

   (a) Use Euler's method with two steps of equal size, starting at $x = -1$, to approximate $f(0)$. Show the work that leads to your answer.

   (b) At the point $(-1, 2)$, the value of $\frac{d^2y}{dx^2}$ is -12. Find the second-degree Taylor polynomial for f about $x = -1$.

   (c) Find the particular solution $y = f(x)$ to the given differential equation with the initial condition $f(-1) = 2$.

WRITE ALL WORK IN THE PINK EXAM BOOKLET.